

TOLUENE, ETHYLBENZENE AND XYLENES: WHAT DO WE KNOW?

When discussing BTEX, benzene rightly attracts particular attention. The other three components – toluene, ethylbenzene and xylenes – nevertheless deserve individual assessment. They share certain physical properties and emission sources but differ in metabolism, toxicity and the importance of different exposure routes. Their potential occurrence in the food chain does not justify assuming that significant quantities are commonly found in food.

THREE SUBSTANCES WITH SIMILAR, BUT NOT IDENTICAL, PROPERTIES

Toluene (methylbenzene) is an aromatic hydrocarbon used as a solvent and found in fuels. Ethylbenzene is used mainly as a feedstock for styrene production and also occurs in petroleum mixtures. Xylenes are three isomers – ortho-, meta- and para-xylene – frequently sold as a mixture. All are volatile organic compounds: they can move from the liquid phase into air, but their actual distribution between air, water, materials and food depends on environmental conditions and matrix properties.

Chemical similarity does not imply a single safety threshold or identical toxicological profiles. Even where several BTEX are measured together, individual results must be distinguished and it must be clear whether the reported xylene concentration refers to one isomer or to their sum.

TOLUENE: SOURCES AND BEHAVIOUR

Well-known sources of toluene include fuels, traffic emissions, paints, adhesives, industrial solvents and processes using petroleum derivatives. Following environmental release it may reach air, water or soil; in surface waters it tends to volatilise and can be degraded by microorganisms. In subsurface systems its persistence depends, among other things, on oxygen availability and local conditions.

For the general population, breathing contaminated air and using certain consumer products are important exposure routes. ATSDR notes that toluene is not frequently detected in food or drinking water. This does not rule out particular contamination incidents, but it precludes describing diet as the dominant route without relevant evidence.

Once absorbed, toluene is mainly metabolised in the liver and eliminated through urinary metabolites and, partly, exhaled air. Effects on the central nervous system are well documented at sufficiently high exposures, especially by inhalation; findings from workplaces or solvent misuse cannot be transferred directly to traces measured in food.

ETHYLBENZENE: EXPOSURE AND UNCERTAINTIES


Ethylbenzene is chiefly associated with petrochemical production and emissions from fuels and solvents. It may contaminate water or soil locally and, in certain conditions, contact raw materials, plant or other materials. Detection in a food sample cannot, by itself, establish the source: information about the supply chain, materials and sampling is required.

Toxicological assessments have considered effects observed in experimental studies, including long-term investigations. Hazard classifications and experimental findings must be read in context; they do not automatically quantify the risk from ingesting low concentrations. When discussing carcinogenicity, it is also essential to identify the agency making the classification and the routes or exposure conditions studied.

XYLENES: A FAMILY OF ISOMERS

'Xylenes' refers to three molecules with the same chemical formula but different positions of their methyl groups on the aromatic ring. They are used industrially and occur in fuels and solvent mixtures. Their volatility favours transfer to air, although contaminated groundwater or production environments may warrant investigation in individual cases.

Critical effects considered in toxicological assessments include those on the nervous system, alongside other effects found experimentally. In food, a physicochemical possibility of transfer must still be distinguished from actually measured occurrence, particularly because exposure studies often focus on air or water rather than the whole diet.



FROM THE ENVIRONMENT TO FOOD: WHICH PATHWAYS ARE PLAUSIBLE?

Water used as an ingredient or in production may act as a carrier if contaminated. Contact with industrial air, improperly used solvents, contaminated materials or surfaces, and accidental events is also possible. Actual significance, however, changes substantially with contact duration, temperature, lipid content, ventilation, packaging and food processing.

It is incorrect to suggest that fatty foods are always the most contaminated: affinity for organic phases is only one factor. Volatilisation during preparation, initial concentration, raw-material origin and storage conditions can all change the outcome. Without comparable surveys, there is no reliable ranking of foods 'most at risk' from toluene, ethylbenzene and xylenes.

WHAT DO DRINKING-WATER GUIDELINE VALUES TELL US?

The World Health Organization has published fact sheets on these substances within its drinking-water quality guidelines. It gives a health-based guideline value of 0.7 mg/L for toluene and 0.5 mg/L for xylenes considered together. These values are based on toxicological and exposure assumptions specific to drinking water; they are not general legal limits for cereals, oil, milk or other foods. The WHO ethylbenzene fact sheet likewise must be interpreted in its precise context, not automatically extended to other matrices.

A health-based guideline value, a legally binding limit and a laboratory limit of quantification are three different concepts. A concentration below the limit of quantification does not necessarily signify absolute absence: it means that, using that method and matrix, reliable quantification below the stated threshold is not possible.

COMBINED EXPOSURE: A LEGITIMATE QUESTION

Several BTEX compounds may coexist at a contaminated site. Assessing a mixture requires information on individual concentrations, exposure routes and potential toxicological interactions. ATSDR has published a specific profile of BTEX mixtures, illustrating why it is wrong either to assume that effects simply add together or that one compound can stand in for all the others.

For food, this question is further complicated by the scarcity of consistent datasets for many matrices. Recognising this limitation is more scientifically defensible than calculating an apparently precise cumulative risk from inadequate evidence.

HOW TO INTERPRET AN ANALYTICAL RESULT

If a laboratory reports toluene, ethylbenzene or xylenes, the initial questions are: which substance or isomer was measured, in which units, and was the result above the limit of quantification? Were containers and procedures used that limit losses and contamination during collection, transport and analysis? Is a control sample available?

Assessing health significance then requires estimating how much of the substance might be ingested, how often and by whom. Analytical detection does not itself imply harm, while the absence of a matrix-specific limit does not establish harmlessness. The analytical methods chapter examines the precautions needed to measure such volatile substances correctly.

IN SUMMARY

Toluene, ethylbenzene and xylenes have distinct industrial uses, environmental sources and toxicological profiles. They may be relevant to targeted investigations of contaminated supply chains or environments, but available information does not support generalised alarm about their routine occurrence in food. A sound approach combines reliable analysis, source identification and realistic exposure assessment, keeping hazard and risk distinct.

SCIENTIFIC AND REGULATORY REFERENCES

- ATSDR (2017). Toxicological Profile for Toluene. [Source](#)
- ATSDR (2004). Interaction Profile for Benzene, Toluene, Ethylbenzene and Xylenes (BTEX). [Source](#)
- ATSDR. Toluene – Toxicological Profile, exposure and environmental fate. [Source](#)
- WHO (2022). Chemical fact sheets: Toluene. [Source](#)
- WHO. Guidelines for Drinking-water Quality – Xylenes, chemical fact sheet. [Source](#)
- WHO (2022). Chemical fact sheets: Ethylbenzene. [Source](#)